

## Substance use in the past 12 months: Lexington, KY Survey dates: June 24–26, 2022 N=227

Page 1 of 2


Lexington, KY

Alcohol

Recreational marijuana

The National Drug Early Warning System (NDEWS) Rapid Street Reporting (RSR) team visited Lexington, KY on June 24-26, 2022. The RSR team conducted surveys (N=227) in public spaces, including parks, nightlife, town squares, sidewalks, malls, and libraries.

For additional context regarding substance use in Kentucky, the Kentucky Injury Prevention and Research Center provided information on state-wide trends. Among the 2,251 fatal drug overdose deaths among Kentucky residents in 2021, 79.6% involved at least one opioid, while 51% involved at least one stimulant. Since 2020, the number of overdose deaths from opioids increased by 12% and 38.8% for stimulants. White individuals accounted for 90.7% of all overdose deaths, while 65.2% of all overdose deaths were among males. Fentanyl was involved in 69% of all drug overdose deaths, followed by methamphetamine at 38.1%.




- + Nonmedical use.
- § Prescription opioids include Vicodin/other hydrocodone, Oxycontin, other oxycodone, Tramadol, codeine, morphine, Dilaudid, methadone, and other prescription opioids.
- \* Prescription stimulants include nonmedical use of Adderall, Ritalin, or another drug for ADHD.
- ‡ Benzodiazepines include Xanax/alpraxolam, Valium/diazepam, Klonopin/clonazopam, and Ativan/lorazepam.
- ${\rm **Other\ opioids\ include\ heroin,\ other\ fentanyls\ such\ as\ carfentanil,\ and\ U-47700/U4.}$
- $\P$  Synthetic cathinones include methodrone, other bath salts, and unknown bath salts.
- △ Other psychedelics include 1P-LSD and ALD-52/Orange Sunshine.



## Substance use in the past 12 months: Lexington, KY Survey dates: June 24–26, 2022 N=227

Page 2 of 2

## **Demographics of Lexington, KY Participants**



7.9% of participants identified as Hispanic/Latino

