


Substance use in the past 12 months: San Diego, CA

Survey dates: August 19–21, 2022
N=391



The National Drug Early Warning System (NDEWS) Rapid Street Reporting (RSR) team visited San Diego, CA on August 19–21, 2022. The RSR team conducted surveys (N=391) in public spaces, including parks, grocery stores, stadium/arenas, libraries, and sidewalks. According to the California Overdose Surveillance dashboard, the rate of drug overdose deaths increased from 21.3 to 26.1 per 100,000 San Diego residents from 2020 to 2021.



† Nonmedical use.

§ Prescription opioids include Vicodin, Oxycontin, Tramadol, codeine, morphine, Dilaudid, methadone, Opana, Suboxone, and other oxycodone.

* Prescription stimulants include Adderall, Ritalin, or another drug for ADHD.

† Benzodiazepines include Xanax, Valium, Klonopin, Ativan, Rohypnol, and other benzodiazepines.

x Other opioids include U-47700 and other fentanyls such as carfentanil.

^ Other tryptamines include 5-MeO-DiPT, 4-AcO-DMT, 5-MeO-MiPT, AMT, and 5-MeO-DMT

†† NPS benzodiazepines include etizolam and clonazolam.

** Dissociative drugs include 3-MeO-PCP, 4-MeO-PCP, and other dissociative drugs.

Δ Euphoric stimulants include MDA and other euphoric stimulants other than MDMA/ecstasy.


£ 2 Cs include 2C-E and 2C-I.

Substance use in the past 12 months: San Diego, CA



Survey dates: August 19–21, 2022
N=391

Demographics of San Diego, CA Participants


Race



Age




Gender identity



23% of participants identified as Hispanic/ Latino

Years of education

Mean years of education: 13.5 years



Location of survey

*Each site type included multiple venues. "Other" includes a grocery store and a library.

