




RAPID STREET REPORTING

Substance use in the past 12 months: San Diego, CA Survey dates: August 19–21, 2022 N=391



The National Drug Early Warning System (NDEWS) Rapid Street Reporting (RSR) team visited San Diego, CA on August 19–21, 2022. The RSR team conducted surveys (N=391) in public spaces, including parks, grocery stores, stadium/arenas, libraries, and sidewalks. According to the California Overdose Surveillance dashboard, the rate of drug overdose deaths increased from 21.3 to 26.1 per 100,000 San Diego residents from 2020 to 2021.



† Nonmedical use.
 § Prescription opioids include Vicodin, Oxycontin, Tramadol, codeine, morphine, Dilaudid, methadone, Opana, Suboxone, and other oxycodone.
 * Prescription stimulants include Adderall, Ritalin, or another drug for ADHD.
 ‡ Benzodiazepines include Xanax, Valium, Klonopin, Ativan, Rohypnol, and other benzodiazepines.
 X Other opioids include U-47700 and other fentanyls such as carfentanil.
 ^ Other tryptamines include 5-MeO-DiPT, 4-AcO-DMT, 5-MeO-MiPT, AMT, and 5-MeO-DMT
 ¶ NPS benzodiazepines include etizolam and clonazolam.
 ** Dissociative drugs include 3-MeO-PCP, 4-MeO-PCP, and other dissociative drugs.
 Δ Euphoric stimulants include MDA and other euphoric stimulants other than MDMA/ecstasy.
 £ 2 Cs include 2C-E and 2C-1.




RAPID STREET REPORTING

Substance use in the past 12 months: San Diego, CA Survey dates: August 19–21, 2022 N=391


Demographics of San Diego, CA Participants

Race



- White
- American Indian
- Black
- Asian
- Other
- Pacific Islander


Age



- 18-29
- 30-39
- 40-49
- 50-59
- 60+

Mean age: 44.5 years

Gender identity




- Male
- Female
- Trans female
- Non-conforming

23% of participants identified as Hispanic/ Latino

Years of education

Mean years of education: 13.5 years



Location of survey

*Each site type included multiple venues. "Other" includes a grocery store and a library.

